

(11) (A) No. 1 234 118

(45) ISSUED R80315

(52) CLASS 260-290.3
C.R. CL. 260-277.9,
 260-296

(51) INT. CL. ⁴ C07D 213/68, 213/89,
 401/12

(19) (CA) **CANADIAN PATENT** (12)

(54) Pyridyl-N-Oxide Intermediates for the Preparation of
Omeprazole

(72) Brandstrom, Arne E.)
Lamm, Bo R.,
Sweden

(73) Granted to Aktiebolaget Hässle
Sweden

(21) APPLICATION No. 431,863
(22) FILED 830706
(30) PRIORITY DATE Sweden (8204879-4) 820826

No. OF CLAIMS 13 - NO DRAWING

Canada

DEPOSITED BY THE PATENT OFFICE, OTTAWA
CCA-274 (11-82)


Best Available Copy

DESCRIPTIONField of the invention

The present invention relates to novel chemical intermediates, a process for their preparation, and their use in the preparation of pharmacologically active substances.


Background of the invention

Compounds of the general formula (i) wherein R^1 and R^2 are the same or different and are each selected from the group consisting of hydrogen, alkyl, halogen, carbomethoxy, alkoxy and alkanoyl have been disclosed in e.g. European patent No. 0005 129 as useful therapeutical compounds. One of these compounds, known under the generic name omeprazole ($R^1 = 5-OCH_3$, $R^2 = H$)



is being developed as a gastric acid secretion inhibiting drug. It can also be used for providing gastrointestinal cytoprotective effects in mammals and man.

It is important to obtain simple and efficient intermediates and routes of synthesis for omeprazole and, in a more general sense, for therapeutically active compounds such as benzimidazole derivatives containing the pyridylmethyl moiety



1234118

2

The present invention provides novel compounds which are useful as intermediates in the preparation of therapeutically active compounds such as benzimidazole derivatives which contain a pyridylmethyl radical of the formula (I), and methods for the preparation of such compounds.


5

Prior art

Substituted benzimidazoles containing a pyridine radical of the formula (I) are disclosed i.a. in European patent 0005 129. A problem with these compounds is their stability characteristics. Upon storage without any special precautions being taken, they are degraded at a rate which is higher than desired. E.g. by storage of omeprazole, which is a substituted benzimidazole disclosed in the patent cited above, at accelerated conditions, that is at +37°C and at a relative humidity of 80% for a period of 6 months, about 6% of the substance is converted to degradation products.

Detailed description of the invention

20 It has been found according to the present invention that the compounds of the formula



25

I

wherein R is H or CH₃, are novel and useful intermediates in the preparation of pharmaceutically useful compounds, e.g. substituted benzimidazoles of the general formula (I). The compounds of the formula I are the products obtained from the preceding nitration reaction (see preparation below), for which the N-oxide form may be considered necessary, and the following substitution reaction in which the pyridine N-oxide form is very advantageous considering the yields.


35

1234118

3

In addition, the N-oxide state of the compounds of the formula I is very advantageous for the subsequent conversion to the 2-hydroxymethylpyridine (procedures A and B). Direct hydroxymethylation of the corresponding non-oxidized pyridines

5



10


only gives low yields (<20%).

B.

The compounds of the formula I may ^{advantageously} ~~advantageously~~ be prepared by processing both the nitration step and the substitution step without isolation of the intermediate nitro-pyridine. Furthermore they are stable and can be stored in bulk form. For example, the compounds according to the invention of the formula I are useful as intermediates in the preparation of the corresponding 2-hydroxymethylpyridine and reactive derivatives thereof of the formula

15

20




25

(iii)

or a salt thereof, in which formula Z is a hydroxy group or reactive esterified hydroxy group, e.g. halogen such as Cl and p-toluenesulfonyl used for the preparation of e.g. omeprazole. The reactive intermediate of the formula (iii) is then reacted in known manner with a benzimidazole derivative of the formula

30

35




(iv)

1234118

4

whereafter oxidation in known manner of the reaction product of the formula




10 yields omeprazole. A preferable method of preparing omeprazole is to use a compound with the general formula I, wherein R is H as an intermediate. The most preferable method of preparing omeprazole is to use a compound, wherein R is CH₃ as an intermediate.

15 The present invention also relates to a process for the preparation of the compounds of the formula I.

20 The compounds of the invention of the formula I are prepared according to the invention by

a) reacting a compound of the formula



30 wherein R is H or CH₃,
with a nitrating agent such as nitric acid




III

35 to the formation of a compound of the formula

1234118

5




IV

5

wherein R has the meaning given above whereafter

- 10 b) the compound of the formula IV is directly reacted with methoxide to give the desired end product of the formula



I

15

wherein R is H or CH₃.

20

The reaction conditions for the steps a) and b) are suitably the following.

- 25 For reaction a), ordinary nitration conditions, i.e., a mixture of conc. sulfuric acid and nitric acid of different concentrations are used. Mixtures containing organic solvents such as acetic acid and nitromethane may also be used.


- 30 For reaction b) a solution of methoxide anion in methanol is preferably used. Methoxide salts in inert solvents such as toluene may also be used. A solution of methoxide in methanol can be prepared from sodium hydroxide and methanol.

- 35 The utilization of the compounds I in the preparation of reactive derivatives of corresponding 2-hydroxymethylpyridine can be carried out as illustrated below;


1234118


6

A. Procedure useful for the preparation of a compound of the formula (iii) utilizing a compound of the formula I wherein R is CH₃:



25 B. Procedure useful for the preparation of a compound of the formula (iii) utilizing a compound of the formula I wherein R is H:





Suitable sources of free radicals are e.g. $(\text{NH}_4)_2\text{S}_2\text{O}_8$ or other salts of persulfuric acid.

10

The compound of the formula (iii) thus obtained, or a salt thereof, is thereafter in known manner as described in the prior art reacted with the desired benzimidazole derivative (iv) as described above.

15 The invention is illustrated by the following examples.

Example 1. Preparation of 2,3,5-trimethyl-4-methoxypyridine-N-oxide

2,3,5-trimethyl-pyridine-N-oxide (1457 g, 10 moles) was dissolved in conc. H_2SO_4 (1200 ml, 22.08 moles) in a 50 litres reaction vessel. A
 20 nitration solution (1750 ml, 32.2 moles conc. H_2SO_4 and 2065 ml, 29.84 moles 65% HNO_3) was added at 90°C during 1 hour. The solution was stirred at 90° for 1.5 hours and thereafter cooled to 30°C. The pH of the reaction mixture was then adjusted by adding 10M NaOH (11.65 litres, 116.5 moles) during cooling with water so that the temperature was
 25 kept below 40°C. The NaOH was added during about 2 hours. Thereafter CH_2Cl_2 (25 litres) was added and the mixture stirred vigorously for 30 minutes. The phases formed were separated and the CH_2Cl_2 -phase was transferred to a 100 litres reaction vessel. The water phase was discarded. The methylenechloride was distilled off. To the remainder was
 30 added 15 l of toluene which was then distilled off under reduced pressure, followed by another 15 l portion of toluene which was also removed by distillation. 8 litres of methanol was added and the mixture heated to boiling temperature. A solution of NaOH (595 g, 14.9 moles) in CH_3OH (16 litres) was added during about 1.5 hours. The reaction mixture
 35 obtained was cooled and its pH adjusted to 8 using conc. H_2SO_4 (250 ml, 4.6 moles). Remaining methanol was distilled off and CH_2Cl_2 (20 litres) was added to the remainder. The mixture was stirred for about 30 minutes and inorganic salts were filtered off and washed with CH_2Cl_2 . The fil-


1234118

8

trates obtained were pooled and evaporated, yielding 1287 g of 2,3,5-trimethyl-4-methoxy-pyridine-N-oxide with a purity of 89%. The identity of the reaction product was confirmed with ^1H and ^{13}C NMR. ^1H -NMR: δ (COCl_2) 2.22(s,3H), 2.27(s,3H), 2.51(s,3H), 3.81(s,3H), 3.18(s,1H).

5

The reaction sequence is:



15 The 2,3,5-trimethylpyridine-N-oxide used as starting material was prepared as follows.

Preparation of 2,3,5-trimethyl-pyridine-N-oxide.

20 To a 100 litres reaction vessel was added 2,3,5-trimethyl-pyridine (10.9 kg, 89.2 moles) and acetic acid (30 litres). The temperature was raised to 90°C. The mixture was stirred for 3 hours and thereafter cooled to 60°C, whereafter H_2O_2 (35% solution, 3122 ml, 35.67 moles) was added during 1 hour. The temperature was then raised to 90°C. The reaction mixture was stirred overnight. After cooling to 40°C an additional amount of H_2O_2 solution (936 ml, 10.7 moles) was added during 1 hour. The temperature was then raised to 90°C. The reaction mixture was stirred for 3 hours and was allowed to stand without heating overnight. Excess of acetic acid was distilled off under vacuum. To the remainder was added NaOH (10M) until pH 10. CH_2Cl_2 (10 litres) was added and the resulting mixture was stirred vigorously. The CH_2Cl_2 phase was separated and the water phase was extracted twice with CH_2Cl_2 (10 litres). The combined CH_2Cl_2 - phases were dried over MgSO_4 and filtered. The filtrate was evaporated yielding 2,3,5-trimethyl-pyridine-N-oxide (11920 g, 94% purity). The identity of the product was confirmed with ^1H and ^{13}C NMR.

35

Example 2. Preparation of 3,5-dimethyl-4-methoxy-pyridine-N-oxide.

- 3,5-dimethyl-pyridine-N-oxide (3500 g, 28.5 moles) was dissolved in conc. H_2SO_4 (3500 ml, 64.4 moles). The solution was cooled to 90°C and nitration solution (5 l, 91.5 moles, conc. H_2SO_4 and 5.9 l, 85 moles 5 65% HNO_3) was added during 4 hours at 90°C. The solution was stirred at 90°C over night. The solution was cooled to 30°C and neutralized with 10M NaOH (36 l, 360 moles) during 4 hours and the temperature kept below 30°C. Acetonitrile (35 litres) was added and the mixture stirred vigorously for 30 minutes. The acetonitrile layer was separated. The 10 extraction procedure was repeated with 15 l of acetonitrile, and the combined acetonitrile were extracted with water (10 l at 60°C). The upper layer was collected and evaporated at reduced pressure (bp 30-55°C/130 mm Hg). Toluene (10 l) was added and remaining water was thoroughly removed by azeotropic distillation at reduced pressure (bp 15 55-65°C/130 mm Hg). Methylalcohol (7 l, 173 moles) was added and the mixture was heated to reflux temperature. A solution of NaOH (1138 g, 28.45 moles) in 30 litres methylalcohol was added over a period of 15 hours. The reaction mixture was cooled and pH adjusted to 9 using conc. HCl (1200 ml, 14 moles). Remaining methanol was evaporated. The 20 residue was cooled and CH_2Cl_2 (30 l) and activated carbon (50 g) were added. The mixture was stirred for 30 minutes, filtered and the residue washed with CH_2Cl_2 . The filtrates were evaporated. The solid product was washed with petroleum ether, (5 litres bp 60-80°C) at 50°C for 30 minutes and filtered. This procedure was repeated once. The product 25 was dried at reduced pressure. Yield 2400 g 3,5-dimethyl-4-methoxypyridine-N-oxide with a purity of 90%. The identity of the product was confirmed with 1H - and ^{13}C -NMR. 1H -NMR: δ ($COCl_3$) 2.23(s,6H), 3.81(2,3H), 8.03(s,2H).
- 30 The 3,5-dimethyl-pyridine-N-oxide used as starting material was prepared as follows.

- 3,5-lutidine (15 kg, 140.2 moles) was dissolved in acetic acid (48 l) at 60°C. Hydrogen peroxide (8430 ml, 92 moles) was added during 3 hours. 35 The solution was heated to 90°C and kept at this temperature for 3 hours. The reaction mixture was cooled to 60°C and hydrogen peroxide (3500 ml, 41 moles) was added during 1 hour. The temperature was raised

1234118


10

- to 90°C and kept there for 16 hours. The reaction mixture was evaporated at reduced pressure (70°C 300 mm Hg). The residue (approx 25 litres) was cooled and pH adjusted to 10 with NaOH-solution (23 litres 10M). Acetonitrile (30 litres) was added and the mixture was stirred for 30 minutes.
- 5 The sodiumacetate was separated off and washed with 10 l acetonitrile. The liquid phase was evaporated at reduced pressure (55°C, 200 mm Hg). The remaining solution (approx 25 litres) was extracted with CH_2Cl_2 (20 litres and 3 x 5 litres). The combined organic layers were dried over MgSO_4 , filtrated and evaporated at reduced pressure (50°C 200 mm
- 10 Hg). When all CH_2Cl_2 had distilled off unreacted 3,5-lutidine was evaporated at 75°C, 8 mm Hg. Yield 14940 g of 3,5-dimethylpyridine-N-oxide. The identity was confirmed with ^1H and ^{13}C NMR.

- The conversion of the compounds of the formula I to 3,5-dimethyl-4-methoxy-2-hydroxymethylpyridine can be carried out according to Procedure A and Procedure B as described above and exemplified below.

Procedure A:

- 20 step 1:




- 2,3,5-dimethyl-4-methoxypyridine-N-oxide (1268 g, 6.75 moles) obtained in Example 1, dissolved in acetic acid (740 ml), was added dropwise to $(\text{CH}_3\text{CO})_2\text{O}$ (2140 ml) heated to 90°C. The heating was discontinued during the addition. The temperature rose to 130°C. Thereafter the reaction solution was stirred for 1 hour and then cooled to 80°C where-
- 30 after CH_3OH (2460 ml) was added. The reaction solution was evaporated and the remainder used directly in step 2.

35


1234118

11

step 2:



To the remainder from step 1 was added NaOH (3300 ml, 10M). The mixture
 10 was refluxed for 5 hours, cooled and extracted with CH_2Cl_2 (8 litres).
 The phases were separated and the water phase extracted with CH_2Cl_2
 (2 x 4 litres). The combined CH_2Cl_2 - phases were dried over MgSO_4 ,
 refluxed with a few grams of decolorizing carbon and filtrated, yielding
 3,5-dimethyl-4-methoxy-2-hydroxy-methylpyridine (941 g). The identity
 15 of the product was confirmed with ^1H and ^{13}C NMR.

Procedure B:

30 3,5-Dimethyl-4-methoxypyridine-N-oxide (61.2 g) obtained in Example 2
 was dissolved in CH_3OH (458 ml). Dimethylsulfate (38 ml 0.4 moles)
 was added dropwise during 15 minutes and pH adjusted to 5.0 using 10M
 NaOH. The mixture was stirred for 15 minutes and thereafter refluxed
 35 for 1 hour. An additional amount of dimethylsulfate (3.8 ml, 0.04 moles)
 was added dropwise and the mixture was refluxed for 1.5 hours. Stirring

1234118


12

was continued overnight at room temperature. Thereafter the mixture was heated to reflux and $(\text{NH}_4)_2\text{C}_2\text{O}_8$ (91.2 g, 0.4 mole) dissolved in water (169 ml) was added during 1.75 hours, followed by refluxing for 1.5 hours and stirring at room temperature overnight. Thereafter CH_3OH (452 ml) was added. Precipitated salts were filtered off and discarded. After evaporation of CH_3OH , the remaining water phase (pH 0.6) was adjusted to pH 10.0 using 10M NaOH (145 ml). The water phase was extracted three times with CH_2Cl_2 . The combined CH_2Cl_2 phases were dried over Na_2SO_4 , evaporated and dried, yielding 3,5-dimethyl-4-methoxy-2-hydroxymethylpyridine (44.2 g). The identity of the product was confirmed with ^1H and ^{13}C NMR and the purity checked with gas chromatography.

1234118


THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE
PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A process for the preparation of a compound of the
formula




I

wherein R is hydrogen or methyl, which process comprises nitrating
a compound of the formula



II

to form a compound of the formula




IV

in which formulas R is hydrogen or methyl, and reacting the compound
of the formula IV thus obtained with a methoxide to give a compound
of the formula

13

B.

1234118



I

in which formulas R is hydrogen or methyl.


2. A process according to claim 1 wherein the reaction with methoxide is carried out in methanol.
3. A process according to claim 1 wherein the nitration is carried out by reaction with nitric acid.
4. A process according to claim 3 wherein the nitric acid is in admixture with sulphuric acid.
5. A process according to claim 1 wherein R is methyl.
6. A process according to claim 1 wherein R is hydrogen.
7. A process according to claim 1 wherein R is methyl and the obtained compound of formula I is subjected to reaction with acetic anhydride followed by alkali to convert the methyl group in the 2-position into a hydroxymethyl group.
8. A process according to claim 1 wherein R is hydrogen and the obtained compound of formula I is subjected to reaction with dimethyl sulphate followed by reaction with methanol in the presence of a source of free radicals to insert in the 2-position a hydroxymethyl group.

14


1234118

23940-447


9. A process according to claim 7 or 8 which comprises the further step of reacting the 2-hydroxymethyl compound with a chlorinating agent to obtain a compound of formula (iii)



10. A process according to claim 7 or 8 which comprises the further step of reacting the 2-hydroxymethyl compound with a chlorinating agent to obtain a compound of formula (iii)



followed by reaction with a benzimidazole derivative of the formula (iv)



and oxidation to yield omeprazole.

11. A compound of formula I as defined in claim 1.
12. A compound of formula I as defined in claim 1 wherein R is methyl.

15

C₁

1234118

23940-447

13. A compound of formula I as defined in claim 1 wherein R is hydrogen.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA

PATENT AGENTS

16

C₁

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.